

Jerzy Trzeciak

Revised edition

# Writing Mathematical Papers in English

a practical guide



European Mathematical Society

Jerzy Trzeciak

Revised edition

# Writing Mathematical Papers in English

a practical guide



European Mathematical Society

Author:

Jerzy Trzeciak  
Publications Department  
Institute of Mathematics  
Polish Academy of Sciences  
00-956 Warszawa  
Poland

Bibliographic information published by Die Deutsche Bibliothek

Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie;  
detailed bibliographic data are available at <http://dnb.ddb.de>.

ISBN 3-03719-014-0

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, recitation, broadcasting, reproduction on microfilms or in other ways, and storage in data banks. For any kind of use permission of the copyright owner must be obtained.

Licensed Edition published by the European Mathematical Society

Contact address:

European Mathematical Society Publishing House  
Seminar for Applied Mathematics  
ETH-Zentrum FLI C4  
CH-8092 Zürich  
Switzerland

Phone: +41 (0)1 632 34 36  
Email: [info@ems-ph.org](mailto:info@ems-ph.org)  
Homepage: [www.ems-ph.org](http://www.ems-ph.org)

First published by Gdańskie Wydawnictwo Oświatowe, ul. Grunwaldzka 413, 80-307 Gdańsk,  
Poland; [www.gwo.pl](http://www.gwo.pl).

© Copyright by Gdańskie Wydawnictwo Oświatowe, 1995

Printed in Germany

9 8 7 6 5 4 3 2 1

## PREFACE

The booklet is intended to provide practical help for authors of mathematical papers. It is written mainly for non-English speaking writers but should prove useful even to native speakers of English who are beginning their mathematical writing and may not yet have developed a command of the structure of mathematical **discourse**.

The booklet is oriented mainly to research mathematics but applies to almost all mathematics writing, except more elementary texts where good teaching praxis **typically** favours substantial repetition and **redundancy**.

There is no intention whatsoever to impose any uniformity of mathematical style. Quite the contrary, the aim is to encourage prospective authors to write structurally correct manuscripts as expressively and flexibly as possible, but without compromising certain basic and universal rules.

The first part provides a collection of ready-made sentences and expressions that most commonly occur in mathematical papers. The examples are divided into sections according to their use (in introductions, definitions, theorems, proofs, comments, references to the literature, acknowledgements, editorial correspondence and referees' reports). Typical errors are also pointed out.

The second part concerns selected problems of English grammar and usage, most often encountered by mathematical writers. Just as in the first part, an abundance of examples are presented, all of them taken from actual mathematical texts.

The author is grateful to Edwin F. Beschler, Daniel Davies, Zofia Denkowska, Zbigniew Lipecki and Zdzisław Skupień for their helpful criticism. Thanks are also due to Adam Mysior and Marcin Adamski for suggesting several improvements, and to Henryka Walas for her painstaking job of typesetting the continuously varying manuscript.

*Jerzy Trzeciak*

# CONTENTS

## Part A: Phrases Used in Mathematical Texts

Abstract and introduction.....	4
Definition.....	6
Notation.....	7
Property.....	8
Assumption, condition, convention.....	10
Theorem: general remarks.....	12
Theorem: introductory phrase.....	13
Theorem: formulation.....	13
Proof: beginning.....	14
Proof: arguments.....	15
Proof: consecutive steps.....	16
Proof: “it is sufficient to ....”.....	17
Proof: “it is easily seen that ....”.....	18
Proof: conclusion and remarks.....	18
References to the literature.....	19
Acknowledgments.....	20
How to shorten the paper.....	20
Editorial correspondence.....	21
Referee’s report.....	21

## Part B: Selected Problems of English Grammar

Indefinite article (a, an, —).....	23
Definite article (the).....	24
Article omission.....	25
Infinitive.....	27
Ing-form.....	29
Passive voice.....	31
Quantifiers.....	32
Number, quantity, size.....	34
How to avoid repetition.....	38
Word order.....	40
Where to insert a comma.....	44
Hyphenation.....	46
Some typical errors.....	46
Index.....	49

# PART A: PHRASES USED IN MATHEMATICAL TEXTS

## ABSTRACT AND INTRODUCTION

We prove that in some families of compacta there are no universal elements.

It is also shown that .....

Some relevant counterexamples are indicated.

It is of interest to know whether ..... We wish to investigate .....

We are interested in finding ..... Our purpose is to .....

It is natural to try to relate ..... to .....

This work was intended as an attempt to motivate (at motivating) .....

The aim of this paper is to bring together two areas in which .....

In		Section 3 the third section [Note: paragraph ≠ section]		we		review some of the standard facts on .....
						have compiled some basic facts .....
						summarize without proofs the relevant material on .....
						give a brief exposition of .....
						briefly sketch .....
						set up notation and terminology.
						discuss (study/treat/examine) the case .....
						introduce the notion of .....
						develop the theory of .....
						will look more closely at .....
						will be concerned with .....
						proceed with the study of .....
						indicate how these techniques may be used to .....
						extend the results of ..... to .....
						derive an interesting formula for .....
						it is shown that .....
						some of the recent results are reviewed in a more general setting.
						some applications are indicated.
						our main results are stated and proved.

Section 4		contains a brief summary (a discussion) of .....
		deals with (discusses) the case .....
		is intended to motivate our investigation of .....
		is devoted to the study of .....
		provides a detailed exposition of .....
		establishes the relation between .....
		presents some preliminaries.

We will		touch only a few aspects of the theory.
		restrict our attention (the discussion/ourselves) to .....

It is not our purpose to study .....

No attempt has been made here to develop .....

It is possible that ..... but we will not develop this point here.

A more complete theory may be obtained by .....

However, | this topic exceeds the scope of this paper.  
| we will not use this fact in any essential way.

The basic (main) | idea is to apply .....  
| geometric ingredient is .....

The crucial fact is that the norm satisfies .....

Our proof involves looking at .....

The proof is | based on the concept of .....  
| similar in spirit to .....  
| adapted from .....

This idea goes back at least as far as [7].

We emphasize that .....

It is worth pointing out that .....

The important point to note here is the form of .....

The advantage of using ..... lies in the fact that .....

The estimate we obtain in the course of proof seems to be of independent interest.

Our theorem provides a natural and intrinsic characterization of .....

Our proof makes no appeal to .....

Our viewpoint sheds some new light on .....

Our example demonstrates rather strikingly that .....

The choice of ..... seems to be the best adapted to our theory.

The problem is that .....

The main difficulty in carrying out this construction is that .....

In this case the method of ..... breaks down.

This class is not well adapted to .....

Pointwise convergence presents a more delicate problem.

The results of this paper were announced without proofs in [8].

The detailed proofs will appear in [8] (elsewhere/in a forthcoming publication).

For the proofs we refer the reader to [6].

It is to be expected that .....

One may conjecture that .....

One may ask whether this is still true if .....

One question still unanswered is whether .....

The affirmative solution would allow one to .....

It would be desirable to ..... but we have not been able to do this.

These results are far from being conclusive.

This question is at present far from being solved.

Our method has the **disadvantage** of not being **intrinsic**.  
 The solution falls short of providing an explicit formula.  
 What is still **lacking** is an explicit description of .....

As for **prerequisites**, the reader is expected to be familiar with .....  
 The first two chapters of ..... constitute sufficient preparation.  
 No preliminary knowledge of ..... is required.

To facilitate access to the individual topics, the chapters are rendered as self-contained as possible.

For the convenience of the reader we repeat the relevant material from [7] without proofs, thus making our exposition self-contained.

## DEFINITION

A set  $S$  is *dense* if .....

A set  $S$  is called (said to be) *dense* if .....

We call a set *dense* (We say that a set is *dense*) if .....

We call  $m$  the *product measure*. [Note the word order after “we call”.]

The function  $f$  is given (defined) by  $f = \dots$

Let  $f$  be given (defined) by  $f = \dots$

We define  $T$  to be  $AB + CD$ .

This map is defined by  $\left\{ \begin{array}{l} \text{requiring } f \text{ to be constant on .....} \\ \text{the requirement that } f \text{ be constant on .....} \\ \text{[Note the infinitive.]} \\ \text{imposing the following condition: .....} \end{array} \right.$

The *length* of a sequence is, by definition, the number of .....

The *length* of  $T$ , denoted by  $l(T)$ , is defined to be .....

By the *length* of  $T$  we mean .....

Define (Let/Set)  $E = Lf$   $\left\{ \begin{array}{l} \text{, where } f \text{ is .....} \\ \text{we have set } f = \dots \\ \text{, } f \text{ being the solution of .....} \\ \text{with } f \text{ satisfying .....} \end{array} \right.$

We will consider  $\left\{ \begin{array}{l} \text{the behaviour of the family } g \text{ defined as follows.} \\ \text{the height of } g \text{ (to be defined later) and .....} \end{array} \right.$

To measure the growth of  $g$  we make the following definition.

In this way we obtain what  $\left\{ \begin{array}{l} \text{we shall call} \\ \text{will be referred to as} \\ \text{is known as} \end{array} \right\}$  the *P-system*.

Since .....,  $\left\{ \begin{array}{l} \text{the norm of } f \text{ is well defined.} \\ \text{the definition of the norm is unambiguous (makes sense).} \end{array} \right.$



It is immaterial which  $M$  we choose to define  $F$  as long as  $M$  contains  $x$ . This product is independent of which member of  $g$  we choose to define it. It is Proposition 8 that makes this definition allowable.

Our definition agrees  $\left\{ \begin{array}{l} \text{with the one given in [7] if } u \text{ is } \dots \\ \text{with the classical one for } \dots \end{array} \right.$

Note that  $\left\{ \begin{array}{l} \text{this coincides with our previously introduced} \\ \text{terminology if } K \text{ is convex.} \\ \text{this is in agreement with [7] for } \dots \end{array} \right.$

## NOTATION

We will denote by  $Z$   $\left\{ \begin{array}{l} \text{Let us denote by } Z \\ \text{Let } Z \text{ denote} \end{array} \right. \left| \begin{array}{l} \text{the set } \dots \\ \text{the set } \dots \end{array} \right. \quad \text{Write } \langle \text{Let/Set} \rangle f = \dots$   
 $\left. \begin{array}{l} \text{Let us denote by } Z \\ \text{Let } Z \text{ denote} \end{array} \right\} \left[ \text{Not: "Denote } f = \dots" \right]$

The closure of  $A$  will be denoted by  $\text{cl}A$ .

We will use the symbol  $\langle \text{letter} \rangle k$  to denote  $\dots$

We write  $H$  for the value of  $\dots$

We will write the negation of  $p$  as  $\neg p$ .

The notation  $aRb$  means that  $\dots$

Such cycles are called homologous (written  $c \sim c'$ ).

Here  
 Here and subsequently,  
 Throughout the proof,  
 In what follows,  
 From now on,

 $\left. \begin{array}{l} \text{Here} \\ \text{Here and subsequently,} \\ \text{Throughout the proof,} \\ \text{In what follows,} \\ \text{From now on,} \end{array} \right\} K \left\{ \begin{array}{l} \text{denotes} \\ \text{stands for} \end{array} \right. \left| \begin{array}{l} \text{the map } \dots \end{array} \right.$ 

We follow the notation of [8]  $\langle \text{used in [8]} \rangle$ .

Our notation differs  $\langle \text{is slightly different} \rangle$  from that of [8].

Let us introduce the temporary notation  $Ff$  for  $gfg$ .

With the notation  $f = \dots$ ,  
 With this notation,  
 In the notation of [8, Ch. 7]  $\left| \begin{array}{l} \text{we have } \dots \end{array} \right.$

If  $f$  is real, it is customary to write  $\dots$  rather than  $\dots$

For simplicity of notation,  
 To  $\langle \text{simplify/shorten} \rangle$  notation,  
 By abuse of notation,  
 For abbreviation,

 $\left. \begin{array}{l} \text{For simplicity of notation,} \\ \text{To } \langle \text{simplify/shorten} \rangle \text{ notation,} \\ \text{By abuse of notation,} \\ \text{For abbreviation,} \end{array} \right\} \text{we} \left\{ \begin{array}{l} \text{write } f \text{ instead of } \dots \\ \text{use the same letter } f \text{ for } \dots \\ \text{continue to write } f \text{ for } \dots \\ \text{let } f \text{ stand for } \dots \end{array} \right.$ 

We abbreviate  $Faub$  to  $b'$ .

We denote it briefly by  $F$ . [Not: "shortly"]

We write it  $F$  for short  $\langle \text{for brevity} \rangle$ . [Not: "in short"]

The Radon–Nikodym property (RNP for short) implies that  $\dots$

We will write it simply  $x$  when no confusion can arise.

It will cause no confusion if we use the same letter to designate a member of  $A$  and its restriction to  $K$ .

We shall write the above expression as  
 The above expression may be written as  
 We can write (4) in the form

$$t = \dots$$

The Greek indices label components of sections of  $E$ .

**Print terminology:**

The expression in italics (in italic type), in large type, in bold print;  
 in parentheses ( ) (= round brackets),  
 in brackets [ ] (= square brackets),  
 in braces { } (= curly brackets), in angular brackets  $\langle \rangle$ ;  
 within the norm signs

Capital letters = upper case letters; small letters = lower case letters;  
 Gothic (German) letters; script (calligraphic) letters (e.g.  $\mathcal{F}$ ,  $\mathcal{G}$ );  
 special Roman (blackboard bold) letters (e.g.  $\mathbb{R}$ ,  $\mathbb{N}$ )

Dot  $\cdot$ , prime  $'$ , asterisk = star  $*$ , tilde  $\sim$ , bar  $\bar{}$  [over a symbol], hat  $\hat{}$ ,  
 vertical stroke (vertical bar)  $|$ , slash (diagonal stroke/slant)  $/$ ,  
 dash  $—$ , sharp  $\#$

Dotted line  $\dots\dots$ , dashed line  $-----$ , wavy line  $\sim\sim\sim\sim$

**PROPERTY**

The $\langle An \rangle$ element	<p>such that <math>\langle</math>with the property that<math>\rangle</math> .....</p> <p>[Not: “such an element that”]</p> <p>with the following properties: .....</p> <p>satisfying <math>Lf = \dots</math></p> <p>with <math>Nf = 1</math> <math>\langle</math>with coordinates <math>x, y, z</math><math>\rangle</math></p> <p>of norm 1 <math>\langle</math>of the form .....<math>\rangle</math></p> <p>whose norm is .....</p> <p>all of whose subsets are .....</p> <p>by means of which <math>g</math> can be computed</p> <p>for which this is true</p> <p>at which <math>g</math> has a local maximum</p> <p>described by the equations .....</p> <p>given by <math>Lf = \dots</math></p> <p>depending only on .....</p> <p><math>\langle</math>independent of .....<math>\rangle</math></p> <p>not in <math>A</math></p> <p>so small that <math>\langle</math>small enough that<math>\rangle</math> .....</p> <p>as above <math>\langle</math>as in the previous theorem<math>\rangle</math></p> <p>so obtained</p> <p>occurring in the cone condition</p> <p>[Note the double “r”.]</p> <p>guaranteed by the assumption .....</p>
----------------------------------	---

The $\langle \text{An} \rangle$ element	we have just defined we wish to study $\langle \text{we used in Chapter 7} \rangle$ to be defined later [= which will be defined] in question under study $\langle \text{consideration} \rangle$
---	--

....., the constant  $C$  being independent of ..... [= where  $C$  is .....]  
 ....., the supremum being taken over all cubes .....  
 ....., the limit being taken in  $L$ .

....., where $C$	is so chosen that ..... is to be chosen later. is a suitable constant. is a conveniently chosen element of ..... involves the derivatives of ..... ranges over all subsets of ..... may be made arbitrarily small by .....
------------------	--

The operators $A_i$	have $\langle \text{share} \rangle$ many of the properties of ..... have still better smoothness properties. lack $\langle \text{fail to have} \rangle$ the smoothness properties of ..... still have norm 1.
are	not merely symmetric but actually self-adjoint. not necessarily monotone. both symmetric and positive-definite. not continuous, nor do they satisfy (2). [Note the inverse word order after “nor”.] neither symmetric nor positive-definite. only nonnegative rather than strictly positive, as one may have expected. any self-adjoint operators, possibly even unbounded. still $\langle \text{no longer} \rangle$ self-adjoint. not too far from being self-adjoint.

The	preceding theorem indicated set above-mentioned group resulting region required $\langle \text{desired} \rangle$ element	[But adjectival clauses with prepositions come <i>after</i> a noun, e.g. “the group defined in Section 1”.]
-----	--	---

Both  $X$  and  $Y$  are finite.  
 Neither  $X$  nor  $Y$  is finite.  
 Both  $X$  and  $Y$  are countable, but neither is finite.  
 Neither of them is finite. [Note: “Neither” refers to *two* alternatives.]  
 None of the functions  $F_i$  is finite.

The set  $X$  is not finite; nor  $\langle \text{neither} \rangle$  is  $Y$ .

Note that  $X$  is not finite, nor is  $Y$  countable. [Note the inversion.]

We conclude that  $X$  is empty  $\left| \begin{array}{l} \text{; so also is } Y. \\ \text{, but } Y \text{ is not.} \end{array} \right.$

Hence  $X$  belongs to  $Y$   $\left| \begin{array}{l} \text{; and so does } Z. \\ \text{, but } Z \text{ does not.} \end{array} \right.$

## ASSUMPTION, CONDITION, CONVENTION

We will make  $\langle \text{need} \rangle$  the following assumptions: .....

From now on we make the assumption: .....

The following assumption will be needed throughout the paper.

Our basic assumption is the following.

Unless otherwise stated  $\langle \text{Until further notice} \rangle$  we assume that .....

In the remainder of this section we assume  $\langle \text{require} \rangle g$  to be .....

In order to get asymptotic results, it is necessary to put some restrictions on  $f$ .

We shall make two standing assumptions on the maps under consideration.

It is required  $\langle \text{assumed} \rangle$  that .....

The requirement on  $g$  is that .....

....., where  $g$   $\left| \begin{array}{l} \text{is subject to the condition } Lg = 0. \\ \text{satisfies the condition } Lg = 0. \\ \text{is merely required to be positive.} \end{array} \right.$

Let us orient  $M$  by  $\left| \begin{array}{l} \text{the requirement that } g \text{ be positive.} \\ \text{[Note the infinitive.]} \\ \text{requiring } g \text{ to be .....} \\ \text{imposing the condition: .....} \end{array} \right.$

Now, (4) holds  $\left| \begin{array}{l} \text{for } \langle \text{provided/whenever/only in case} \rangle p \neq 1. \\ \text{unless } p = 1. \\ \text{under } \left| \begin{array}{l} \text{the condition } \langle \text{hypothesis} \rangle \text{ that .....} \\ \text{the more general assumption that .....} \\ \text{some further restrictions on .....} \\ \text{additional } \langle \text{weaker} \rangle \text{ assumptions.} \end{array} \right. \end{array} \right.$

It  $\left| \begin{array}{l} \text{satisfies } \langle \text{fails to satisfy} \rangle \text{ the assumptions of .....} \\ \text{has the desired } \langle \text{asserted} \rangle \text{ properties.} \\ \text{provides the desired diffeomorphism.} \\ \text{still satisfies } \langle \text{need not satisfy} \rangle \text{ the requirement that .....} \\ \text{meets this condition.} \\ \text{does not necessarily have this property.} \\ \text{satisfies all the other conditions for membership of } X. \end{array} \right.$

There is no loss of generality in assuming .....

Without loss  $\langle \text{restriction} \rangle$  of generality we can assume .....

This involves no loss of generality.

We can certainly assume that .....  $\left\{ \begin{array}{l} \text{, since otherwise .....} \\ \text{, for ..... [= because]} \\ \text{, for if not, we replace .....} \\ \text{. Indeed, .....} \end{array} \right.$

Neither the hypothesis nor the conclusion is affected if we replace .....

By choosing  $b = a$  we may actually assume that .....

If  $f = 1$ , which we may assume, then .....

For simplicity (convenience) we ignore the dependence of  $F$  on  $g$ .

[E.g. in notation]

It is convenient to choose .....

We can assume, by decreasing  $k$  if necessary, that .....

Thus  $F$  meets  $S$  transversally, say at  $F(0)$ .

There exists a minimal element, say  $n$ , of  $F$ .

Hence  $G$  acts on  $H$  as a multiple (say  $n$ ) of  $V$ .

For definiteness (To be specific), consider .....

This condition  $\left\{ \begin{array}{l} \text{is not particularly restrictive.} \\ \text{is surprisingly mild.} \\ \text{admits (rules out/excludes) elements of .....} \\ \text{is essential to the proof.} \\ \text{cannot be weakened (relaxed/improved/omitted/} \\ \text{dropped).} \end{array} \right.$

The theorem is true if "open" is deleted from the hypotheses.

The assumption ..... is superfluous (redundant/unnecessarily restrictive).

We will now show how to dispense with the assumption on .....

Our lemma does not involve any assumptions about curvature.

We have been working under the assumption that .....

Now suppose that this is no longer so.

To study the general case, take .....

For the general case, set .....

The map  $f$  will be viewed (regarded/thought of) as  $\left\{ \begin{array}{l} \text{a functor .....} \\ \text{realizing .....} \end{array} \right.$

From now on we  $\left\{ \begin{array}{l} \text{think of } L \text{ as being constant.} \\ \text{regard } f \text{ as a map from .....} \\ \text{tacitly assume that .....} \end{array} \right.$

It is understood that  $r \neq 1$ .

We adopt (adhere to) the convention that  $0/0=0$ .

## THEOREM: GENERAL REMARKS

This theorem	<table style="border-collapse: collapse; width: 100%;"> <tr> <td style="vertical-align: middle; padding-right: 5px;">is</td> <td style="border-left: 1px solid black; padding-left: 5px;">                 an extension (a fairly straightforward generalization/a sharpened version/a refinement) of .....                  an analogue of .....                  a reformulation (restatement) of .....                      in terms of .....                  analogous to .....                  a partial converse of .....                  an answer to a question raised by .....                  deals with .....                  ensures the existence of .....                  expresses the equivalence of .....                  provides a criterion for .....                  yields information about .....                  makes it legitimate to apply .....             </td> </tr> </table>	is	an extension (a fairly straightforward generalization/a sharpened version/a refinement) of ..... an analogue of ..... a reformulation (restatement) of ..... in terms of ..... analogous to ..... a partial converse of ..... an answer to a question raised by ..... deals with ..... ensures the existence of ..... expresses the equivalence of ..... provides a criterion for ..... yields information about ..... makes it legitimate to apply .....
is	an extension (a fairly straightforward generalization/a sharpened version/a refinement) of ..... an analogue of ..... a reformulation (restatement) of ..... in terms of ..... analogous to ..... a partial converse of ..... an answer to a question raised by ..... deals with ..... ensures the existence of ..... expresses the equivalence of ..... provides a criterion for ..... yields information about ..... makes it legitimate to apply .....		

The theorem states (asserts/shows) that .....

Roughly (Loosely) speaking, the formula says that .....

When  $f$  is open, (3.7) just amounts to saying that .....  
to the fact that .....

Here is another way of stating (c): .....

Another way of stating (c) is to say: .....

An equivalent formulation of (c) is: .....

Theorems 2 and 3 may be summarized by saying that .....

Assertion (ii) is nothing but the statement that .....

Geometrically speaking, the hypothesis is that .....; part of the conclusion is that .....

The interest The principal significance The point	<table style="border-collapse: collapse; width: 100%;"> <tr> <td style="vertical-align: middle; padding-right: 5px;">of the lemma is</td> <td style="border-left: 1px solid black; padding-left: 5px;">                 in the assertion .....                  that it allows one to .....             </td> </tr> </table>	of the lemma is	in the assertion ..... that it allows one to .....
of the lemma is	in the assertion ..... that it allows one to .....		

The theorem gains in interest if we realize that .....

The theorem	<table style="border-collapse: collapse; width: 100%;"> <tr> <td style="vertical-align: middle; padding-right: 5px;">is still true</td> <td style="border-left: 1px solid black; padding-left: 5px;">if</td> <td style="border-left: 1px solid black; padding-left: 5px;">we drop the assumption .....</td> </tr> <tr> <td style="vertical-align: middle; padding-right: 5px;">still holds</td> <td style="border-left: 1px solid black; padding-left: 5px;"></td> <td style="border-left: 1px solid black; padding-left: 5px;">it is just assumed that .....</td> </tr> </table>	is still true	if	we drop the assumption .....	still holds		it is just assumed that .....
is still true	if	we drop the assumption .....					
still holds		it is just assumed that .....					

If we take $f = \dots$ Replacing $f$ by $-f$ ,	<table style="border-collapse: collapse; width: 100%;"> <tr> <td style="vertical-align: middle; padding-right: 5px;">we recover</td> <td style="border-left: 1px solid black; padding-left: 5px;">the standard lemma .....</td> </tr> <tr> <td style="vertical-align: middle; padding-right: 5px;"></td> <td style="border-left: 1px solid black; padding-left: 5px;">[7, Theorem 5].</td> </tr> </table>	we recover	the standard lemma .....		[7, Theorem 5].
we recover	the standard lemma .....				
	[7, Theorem 5].				

This specializes to the result of [7] if  $f = g$ .

This result will	<table style="border-collapse: collapse; width: 100%;"> <tr> <td style="vertical-align: middle; padding-right: 5px;">be needed in</td> <td style="border-left: 1px solid black; padding-left: 5px;">prove extremely useful in</td> <td style="border-left: 1px solid black; padding-left: 5px;">Section 8.</td> </tr> <tr> <td style="vertical-align: middle; padding-right: 5px;">not be needed until</td> <td style="border-left: 1px solid black; padding-left: 5px;"></td> <td style="border-left: 1px solid black; padding-left: 5px;"></td> </tr> </table>	be needed in	prove extremely useful in	Section 8.	not be needed until		
be needed in	prove extremely useful in	Section 8.					
not be needed until							

## THEOREM: INTRODUCTORY PHRASE

We have thus proved ..... We can now | rephrase Theorem 8 as follows.  
Summarizing, we have ..... | state the analogue of .....  
| formulate our main results.

We are thus led to the following strengthening of Theorem 6: .....  
The remainder of this section will be devoted to the proof of .....

The continuity of  $A$  is established by our next theorem.  
The following result may be proved in much the same way as Theorem 6.  
Here are some elementary properties of these concepts.  
Let us mention two important consequences of the theorem.  
We begin with a general result on such operators.

[*Note:* Sentences of the type “We now have the following lemma”,  
carrying no information, can in general be cancelled.]

## THEOREM: FORMULATION

If ..... and if ....., then .....

Let  $M$  be ..... | Suppose that ..... | Then ....., | provided  $m \neq 1$ .  
| Assume that ..... | unless  $m = 1$ .  
| Write ..... | with  $g$  a constant  
| satisfying .....

Furthermore (Moreover), .....  
In fact, ..... [= To be more precise]  
Accordingly, ..... [= Thus]

Given any  $f \neq 1$  suppose that ..... Then .....

Let  $P$  satisfy | the hypotheses of ..... | Then .....  
| the above assumptions. |  
|  $N(P) = 1$ .

Let assumptions 1–5 hold. Then .....  
Under the above assumptions, .....  
Under the same hypotheses, .....  
Under the conditions stated above, .....  
Under the assumptions of Theorem 2 with “convergent”  
replaced by “weakly convergent”, .....  
Under the hypotheses of Theorem 5, if moreover .....

Equality holds in (8) if and only if .....  
The following conditions are equivalent: .....

[*Note:* Expressions like “the following inequality holds” can in  
general be dropped.]

## PROOF: BEGINNING

We  
Let us | first | prove ⟨show/recall/observe⟩ that .....  
| | prove a reduced form of the theorem.  
| | outline ⟨give the main ideas of⟩ the proof.  
| | examine  $Bf$ .

But  $A = B$ . | To see ⟨prove⟩ this, let  $f = \dots$   
| We prove this as follows.  
| This is proved by writing  $g = \dots$

We first compute  $If$ . | To this end, consider .....  
| [= For this purpose; *not*: “To this aim”]  
| To do this, take .....  
| For this purpose, we set .....

To deduce (3) from (2), take .....

We claim that ..... Indeed, .....

We begin by proving ..... ⟨by recalling the notion of .....⟩

Our proof starts with the observation that .....

The procedure is to find .....

The proof consists in the construction of .....

The proof is | straightforward ⟨quite involved⟩.  
| by induction on  $n$ .  
| left to the reader.  
| based on the following observation.

The main ⟨basic⟩ idea of the proof is to take .....

The proof | falls naturally into three parts.  
| will be divided into three steps.

We have divided the proof into a sequence of lemmas.

Suppose | the assertion of the lemma is false.  
| , contrary to our claim, that .....

Conversely ⟨To obtain a contradiction⟩, | suppose that .....

On the contrary,  
Suppose the lemma were false. Then we could find .....

If | there existed an  $x$  ....., | we would have .....  
|  $x$  were not in  $B$ , | there would be .....  
| it were true that .....,

Assume the formula holds for degree  $k$ ; we will prove it for  $k + 1$ .

Assuming (5) to hold for  $k$ , we will prove it for  $k + 1$ .

We give the proof only for the case  $n = 3$ ; the other cases are left to the reader.

We give only the main ideas of the proof.



## PROOF: ARGUMENTS

By	definition, ..... the definition of ..... assumption, ..... the compactness of ..... Taylor's formula, ..... a similar argument, ..... the above, ..... the lemma below, ..... continuity, .....	But $f = g$        Theorem 4 now	, which follows from ..... as was described ⟨shown/mentioned/ noted⟩ in .....  shows that ..... yields ⟨gives/ implies⟩ $f = \dots$ leads to $f = \dots$
----	--	--	--

Since  $f$  is compact,  $\left\{ \begin{array}{l} Lf = 0. \quad [Not: \text{“Since } \dots, \text{ then } \dots\text{”}] \\ \text{we have } Lf = 0. \\ \text{it follows that } Lf = 0. \\ \text{we see } \langle \text{conclude} \rangle \text{ that } Lf = 0. \end{array} \right.$

But  $Lf = 0$  since  $f$  is compact.

We have  $Lf = 0$ , because ..... [+ a longer explanation]

We must have  $Lf = 0$ , for otherwise we can replace .....

As  $f$  is compact we have  $Lf = 0$ .

Therefore  $Lf = 0$  by Theorem 6.

That  $Lf = 0$  follows from Theorem 6.

From	this (5) what has already been proved,	we conclude ⟨deduce/see⟩ that ..... we have ⟨obtain⟩ $M = N$ . [Note: without “that”] it follows that ..... it may be concluded that .....
------	---	--

According to ⟨On account of⟩ the above remark, we have  $M = N$ .

It follows that  
 Hence ⟨Thus/Consequently,/Therefore⟩  $M = N$ .

[hence = from this; thus = in this way; therefore = for this reason;  
 it follows that = from the above it follows that]

This gives $M = N$ . We thus get $M = N$ . The result is $M = N$ . Now (3) becomes $M = N$ . This clearly forces $M = N$ .	It is compact,	and so $M = N$ . and consequently $M = N$ . and, in consequence, $M = N$ . and hence bounded. which gives ⟨implies/ yields⟩ $M = N$ . [Not: “what gives”]
--	----------------	---

Now  $F = G = H$ ,  $\left\{ \begin{array}{l} \text{the last equality being a consequence of Theorem 7.} \\ \text{which is due to the fact that } \dots \end{array} \right.$

Since ....., (2) shows that ....., by (4).

We conclude from (5) that ....., hence that ....., and finally that .....

The equality  $f = g$ , which is part of the conclusion of Theorem 7, implies that .....

As in the proof of Theorem 8, equation (4) gives .....

Analysis similar to that in the proof of Theorem 5 shows that .....

[*Note*: “similar as in”]

A passage to the limit similar to the above implies that .....

Similarly (Likewise), .....

Similar arguments apply |  
The same reasoning applies | to the case .....

The same conclusion can be drawn for .....

This follows by the same method as in .....

The term  $Tf$  can be handled in much the same way, the only difference being in the analysis of .....

In the same manner we can see that .....

The rest of the proof runs as before.

We now apply this argument again, with  $I$  replaced by  $J$ , to obtain .....

### PROOF: CONSECUTIVE STEPS

Consider .....	Define	$f = \dots$	Let us	evaluate .....
Choose .....	Let			compute .....
Fix .....	Set			apply the formula to .....
				suppose for the moment .....
				regard $s$ as fixed and .....

[*Note*: The imperative mood is used when you *order* the reader to do something, so you should not write e.g. “Give an example of .....” if you mean “We give an example of .....”]

Adding $g$ to the left-hand side		yields (gives) $h = \dots$
Subtracting (3) from (5)		we obtain (get/have) $f = g$
Writing (Taking) $h = Hf$		[ <i>Note</i> : without “that”]
Substituting (4) into (6)		we conclude (deduce/see) that .....
Combining (3) with (6)		we can assert that .....
Combining these		we can rewrite (5) as .....
[E.g. these inequalities]		
Replacing (2) by (3)		
Letting $n \rightarrow \infty$		
Applying (5)		
Interchanging $f$ and $g$		

[*Note*: The ing-form is either the subject of a sentence (“Adding .....

gives”), or requires the subject “we” (“Adding .....

we obtain”); so do *not* write e.g. “Adding .....

the proof is complete.”]

We continue in this fashion obtaining (to obtain)  $f = \dots$

We may now integrate  $k$  times to conclude that .....

Repeated application of Lemma 6 enables us to write .....  
 We now proceed by induction.  
 We can now proceed analogously to the proof of .....

We next 

claim $\langle$ show/prove that $\rangle$ .....	sharpen these results and prove that .....
---	--

Our next 

claim is that .....	goal is to determine the number of .....
	objective is to evaluate the integral $I$ .
	concern will be the behaviour of .....

We now turn to the case  $f \neq 1$ .

We are now in a position to show ..... [= We are able to]

We proceed to show that .....

The task is now to find .....

Having disposed of this preliminary step, we can now return to .....

We wish to arrange that  $f$  be as smooth as possible.

[Note the infinitive.]

We are thus looking for the family .....

We have to construct .....

In order to get this inequality, it 

will be necessary to .....	is convenient to .....
----------------------------	------------------------

To deal with  $If$ ,  
 To estimate the other term, 

we note that .....
--------------------

  
 For the general case,

**PROOF: "IT IS SUFFICIENT TO ....."**

It 

suffices	to	show $\langle$ prove $\rangle$ that .....
is sufficient		make the following observation.
		use (4) together with the observation that .....

We need only consider three cases: .....

We only need to show that .....

It remains to prove that .....  $\langle$ to exclude the case when .....

What is left is to show that .....

We are reduced to proving (4) for .....

We are left with the task of determining .....

The only point remaining concerns the behaviour of .....

The proof is completed by showing that .....

We shall have established the lemma if we prove the following: .....

If we prove that ....., the assertion follows.

The statement  $O(g)=1$  will be proved once we prove the lemma below.

## PROOF: "IT IS EASILY SEEN THAT ....."

It is	clear (evident/immediate/obvious) that .....
	easily seen that .....
	easy to check that .....
	a simple matter to .....

We see (check) at once that ..... (3).  
 They are easily seen to be smooth. ...., as is easy to check.

It follows easily (immediately) that .....

Of course (Clearly/Obviously), .....

The proof is straightforward (standard/immediate).

An easy computation (A trivial verification) shows that .....

(2) makes it obvious that ..... [= By (2) it is obvious that]

The factor  $Gf$  poses no problem because  $G$  is .....

## PROOF: CONCLUSION AND REMARKS

....., which [Not: "what"] This	proves the theorem.
	completes the proof.
	establishes the formula.
	is the desired conclusion.
	is our claim (assertion). [Not: "thesis"]

gives (4) when substituted in (5) (combined with (5)).

....., and	the proof is complete.
	this is precisely the assertion of the lemma.
	the lemma follows.
	(3) is proved. $f = g$ as claimed (required).

This contradicts our assumption (the fact that .....).

....., contrary to (3).

....., which is impossible. [Not: "what is"]

....., which contradicts the maximality of .....

....., a contradiction.

The proof for  $G$  is similar.

The map  $G$  may be handled in much the same way.

Similar considerations apply to  $G$ .

The same proof	works (remains valid) for .....
	obtains (fails) when we drop the assumption .....

The method of proof carries over to domains .....

The proof above gives more, namely  $f$  is .....

A slight change in the proof actually shows that .....

Note that we have actually proved that .....  
[= We have proved more, namely that .....]

We have used  $\left\{ \begin{array}{l} \text{only the fact that .....} \\ \text{the existence of only the right-hand derivative.} \end{array} \right.$

For  $f=1$   $\left\{ \begin{array}{l} \text{it is no longer true that .....} \\ \text{the argument breaks down.} \end{array} \right.$

The proof strongly depended on the assumption that .....

Note that we did not really have to use .....; we could have applied .....

For more details we refer the reader to [7].

The details are left to the reader.

We leave it to the reader to verify that ..... [Note: the “it” is necessary]

This finishes the proof, the detailed verification of (4) being left to the reader.

## REFERENCES TO THE LITERATURE

(see for instance [7, Th. 1]) (see [7] and the references given there)

(see [Ka2] for  $\left\{ \begin{array}{l} \text{more details)} \\ \text{the definition of .....)} \\ \text{the complete bibliography)} \end{array} \right.$

The best general reference here  $\left\{ \begin{array}{l} \text{is .....} \\ \text{The standard work on .....} \\ \text{The classical work here} \end{array} \right.$  This  $\left\{ \begin{array}{l} \text{was proved by Lax [8].} \\ \text{can be found in} \\ \text{Lax [7, Ch. 2].} \end{array} \right.$

This construction  $\left\{ \begin{array}{l} \text{is due to Strang [8].} \\ \text{goes back to the work of .....} \\ \text{as far as [8].} \\ \text{was motivated by [7].} \\ \text{generalizes that of [7].} \\ \text{follows [7].} \\ \text{is adapted from [7] (appears in [7]).} \\ \text{has previously been used by Lax [7].} \end{array} \right.$

For  $\left\{ \begin{array}{l} \text{a recent account of the theory} \\ \text{a treatment of a more general case} \\ \text{a fuller (thorough) treatment} \\ \text{a deeper discussion of .....} \\ \text{direct constructions along more} \\ \text{classical lines} \\ \text{yet another method} \end{array} \right.$  we refer the reader to [7].

We introduce the notion of ....., following Kato [7].

We follow [Ka] in assuming that .....

The main results of this paper were announced in [7].  
Similar results have been obtained independently by Lax and are to be published in [7].

## ACKNOWLEDGMENTS

The author | wishes to express his thanks ⟨gratitude⟩ to .....  
                  | is greatly indebted to .....  
                  | his active interest in the publication of this paper.  
                  | suggesting the problem and for many stimulating conversations.  
for              | several helpful comments concerning .....  
                  | drawing the author's attention to .....  
                  | pointing out a mistake in .....  
                  | his collaboration in proving Lemma 4.

The author gratefully acknowledges the many helpful suggestions of .....  
during the preparation of the paper.  
This is part of the author's Ph.D. thesis, written under the supervision  
of ..... at the University of .....  
The author wishes to thank the University of ....., where the paper was  
written, for financial support ⟨for the invitation and hospitality⟩.

## HOW TO SHORTEN THE PAPER

General rules:

1. Remember: you are writing for an expert. Cross out all that is trivial or routine.
2. Avoid repetition: do not repeat the assumptions of a theorem at the beginning of its proof, or a complicated conclusion at the end of the proof. Do not repeat the assumptions of a previous theorem in the statement of a next one (instead, write e.g. "Under the hypotheses of Theorem 1 with  $f$  replaced by  $g$ , ....."). Do not repeat the same formula—use a label instead.
3. Check all formulas: is each of them necessary?

Phrases you can cross out:

We denote by  $\mathbb{R}$  the set of all real numbers.  
We have the following lemma.  
The following lemma will be useful.  
..... the following inequality is satisfied:

Phrases you can shorten (see also p. 38):

Let  $\varepsilon$  be an arbitrary but fixed positive number  $\rightsquigarrow$  Fix  $\varepsilon > 0$   
Let us fix arbitrarily  $x \in X$   $\rightsquigarrow$  Fix  $x \in X$   
Let us first observe that  $\rightsquigarrow$  First observe that  
We will first compute  $\rightsquigarrow$  We first compute  
Hence we have  $x=1$   $\rightsquigarrow$  Hence  $x=1$   
Hence it follows that  $x=1$   $\rightsquigarrow$  Hence  $x=1$

Taking into account (4)  $\rightsquigarrow$  By (4)

By virtue of (4)  $\rightsquigarrow$  By (4)

By relation (4)  $\rightsquigarrow$  By (4)

In the interval  $[0, 1]$   $\rightsquigarrow$  In  $[0, 1]$

There exists a function  $f \in C(X)$   $\rightsquigarrow$  There exists  $f \in C(X)$

For every point  $p \in M$   $\rightsquigarrow$  For every  $p \in M$

It is defined by the formula  $F(x) = \dots$   $\rightsquigarrow$  It is defined by  $F(x) = \dots$

Theorem 2 and Theorem 5  $\rightsquigarrow$  Theorems 2 and 5

This follows from (4), (5), (6) and (7)  $\rightsquigarrow$  This follows from (4)–(7)

For details see [3], [4] and [5]  $\rightsquigarrow$  For details see [3]–[5]

The derivative with respect to  $t$   $\rightsquigarrow$  The  $t$ -derivative

A function of class  $C^2$   $\rightsquigarrow$  A  $C^2$  function

For arbitrary  $x$   $\rightsquigarrow$  For all  $x$  (For every  $x$ )

In the case  $n = 5$   $\rightsquigarrow$  For  $n = 5$

This leads to a contradiction with the maximality of  $f$

$\rightsquigarrow \dots$ , contrary to the maximality of  $f$

Applying Lemma 1 we conclude that  $\rightsquigarrow$  Lemma 1 shows that

$\dots$ , which completes the proof  $\rightsquigarrow \dots$  ■

## EDITORIAL CORRESPONDENCE

I would like to submit | the enclosed manuscript “.....”  
I am submitting | for publication in *Studia Mathematica*.

I have also included a reprint of my article ..... for the convenience of the referee.

I wish to withdraw my paper ..... as I intend to make a major revision of it.

I regret any inconvenience this may have caused you.

I am very pleased that the paper will appear in *Fundamenta*.

Thank you very much for accepting my paper for publication in .....

Please find enclosed two copies of the revised version.

As the referee suggested, I inserted a reference to the theorem of .....

We have followed the referee’s suggestions.

I have complied with almost all suggestions of the referee.

## REFeree’S REPORT

The author proves the interesting result that .....

The proof is short and simple, and the article well written.

The results presented are original.

The paper is a good piece of work on a subject that attracts considerable attention.

I am pleased to		recommend it for publication in Studia Mathematica.
It is a pleasure to		
I strongly		

The only remark I wish to make is that condition  $B$  should be formulated more carefully.

A few minor typographical errors are listed below.

I have indicated various corrections on the manuscript.

The results obtained are not particularly surprising and will be of limited interest.

The results are		correct but only moderately interesting.
		rather easy modifications of known facts.

The example is worthwhile but not of sufficient interest for a research article.

The English of the paper needs a thorough revision.

The paper does not meet the standards of your journal.

Theorem 2 is false		as stated.
		in this generality.

Lemma 2 is known (see .....)

Accordingly, I recommend that the paper be rejected.



## PART B: SELECTED PROBLEMS OF ENGLISH GRAMMAR

### INDEFINITE ARTICLE (a, an, —)

*Note:* Use “a” or “an” depending on *pronunciation* and not spelling, e.g. a unit, an  $x$ .

#### 1. Instead of the number “one”:

The four centres lie in **a** plane.

**A** chapter will be devoted to the study of expanding maps.

For this, we introduce **an** auxiliary variable  $z$ .

#### 2. Meaning “member of a class of objects”, “some”, “one of”:

Then  $D$  becomes **a** locally convex space with dual space  $D'$ .

The right-hand side of (4) is then **a** bounded function.

This is easily seen to be **an** equivalence relation.

Theorem 7 has been extended to **a** class of boundary value problems.

This property is **a** consequence of the fact that .....

Let us now state **a** corollary of Lebesgue’s theorem for .....

After **a** change of variable in the integral we get .....

We thus obtain the estimate ..... with **a** constant  $C$ .

*in the plural:*

The existence of **partitions** of unity may be proved by .....

The definition of **distributions** implies that .....

....., with suitable constants.

....., where  $G$  and  $F$  are **differential operators**.

#### 3. In definitions of classes of objects

(i.e. when there are many objects with the given property):

**A** fundamental solution is a function satisfying .....

We call  $C$  **a** module of ellipticity.

**A** classical example of **a** constant  $C$  such that .....

We wish to find **a** solution of (6) which is of the form .....

*in the plural:*

The elements of  $D$  are often called **test functions**.

the set of  $\left\{ \begin{array}{l} \text{points with distance 1 from } K \\ \text{all functions with compact support} \end{array} \right.$

The integral may be approximated by **sums** of the form .....

Taking in (4) **functions**  $v$  which vanish in  $U$  we obtain .....

Let  $f$  and  $g$  be **functions** such that .....

4. In the plural—when you are referring to each element of a class:

Direct sums exist in the category of abelian groups.

In particular, closed sets are Borel sets.

Borel measurable functions are often called Borel mappings.

This makes it possible to apply  $H_2$ -results to functions in any  $H_p$ .

*If you are referring to all elements of a class, use “the”:*

**The** real measures form a subclass of **the** complex ones.

5. In front of an adjective which is intended to mean “having this particular quality”:

This map extends to all of  $M$  in **an** obvious fashion.

**A** remarkable feature of the solution should be stressed.

Section 1 | gives **a** condensed exposition of .....  
describes in **a** unified manner the recent results .....

**A** simple computation gives .....

Combining (2) and (3) we obtain, with **a** new constant  $C$ , .....

**A** more general theory must be sought to account for these irregularities.

The equation (3) has **a** unique solution  $g$  for every  $f$ .

*But:* (3) has **the** unique solution  $g = ABf$ .

## DEFINITE ARTICLE (the)

1. Meaning “mentioned earlier”, “that”:

Let  $A \subset X$ . If  $aB = 0$  for every  $B$  intersecting **the** set  $A$ , then .....

Define  $\exp x = \sum x^i/i!$ . **The** series can easily be shown to converge.

2. In front of a noun (possibly preceded by an adjective) referring to a single, uniquely determined object (e.g. in definitions):

Let  $f$  be **the** linear form  $\left| \begin{array}{l} g \mapsto (g, F). \\ \text{defined by (2).} \end{array} \right.$  [If there is only one.]

So  $u = 1$  in **the** compact set  $K$  of all points at distance 1 from  $L$ .

We denote by  $B(X)$  **the** Banach space of all linear operators in  $X$ .

....., under **the** usual boundary conditions.

....., with **the** natural definitions of addition and multiplication.

Using **the** standard inner product we may identify .....

3. In the construction: the + property (or another characteristic) + of + object:

**The** continuity of  $f$  follows from .....

**The** existence of test functions is not evident.

There is a fixed compact set containing **the** supports of all the  $f^j$ .

Then  $x$  is **the** centre of an open ball  $U$ .

**The** intersection of a decreasing family of such sets is convex.

*But:* Every nonempty open set in  $\mathbb{R}^k$  is a **union** of disjoint boxes.  
 [If you wish to stress that it is some union of not too well specified objects.]

4. In front of a cardinal number if it embraces all objects considered:

**The** two groups have been shown to have the same number of generators. [Two groups only were mentioned.]

Each of **the** three products on the right of (4) satisfies .....  
 [There are exactly three products there.]

5. In front of an ordinal number:

**The first** Poisson integral in (4) converges to  $g$ .

**The second** statement follows immediately from **the first**.

6. In front of surnames used attributively:

<b>the</b> Dirichlet problem	<i>But:</i>	Taylor's formula [without "the"] a Banach space
<b>the</b> Taylor expansion		
<b>the</b> Gauss theorem		

7. In front of a noun in the plural if you are referring to a class of objects as a whole, and not to particular members of the class:

**The** real measures form a subclass of **the** complex ones.

This class includes **the** Helson sets.

## ARTICLE OMISSION

1. In front of nouns referring to activities:

Application of Definition 5.9 gives (45).

**Repeated** application (use) of (4.8) shows that .....

The last formula can be derived by **direct** consideration of .....

Thus  $A$  is the smallest possible extension in which **differentiation** is always possible.

Using integration by parts we obtain .....

If we apply **induction** to (4), we get .....

**Addition** of (3) and (4) gives .....

This reduces the solution to **division** by  $Px$ .

**Comparison** of (5) and (6) shows that .....

2. In front of nouns referring to properties if you mention no particular object:

In questions of **uniqueness** one usually has to consider .....

By **continuity**, (2) also holds when  $f = 1$ .

By **duality** we easily obtain the following theorem.

Here we do not require **translation invariance**.

3. After certain expressions with “of”:

a <b>type of</b> convergence	the <b>hypothesis of</b> positivity
a <b>problem of</b> uniqueness	the <b>method of</b> proof
the <b>condition of</b> ellipticity	the <b>point of</b> increase

4. In front of numbered objects:

It follows from **Theorem 7** that .....

**Section 4** gives a concise presentation of .....

**Property (iii)** is called the triangle inequality.

This has been proved in **part (a)** of the proof.

*But:* the set of solutions of **the** form (4.7)

To prove **the** estimate (5.3) we first extend .....

We thus obtain **the** inequality (3). [*Or:* inequality (3)]

**The** asymptotic formula (3.6) follows from .....

Since **the** region (2.9) is in  $U$ , we have .....

5. To avoid repetition:

the order and symbol of a distribution

the associativity and commutativity of  $A$

the direct sum and direct product

the inner and outer factors of  $f$  [Note the plural.]

*But:* a deficit or an excess

6. In front of surnames in the possessive:

Minkowski's inequality, *but:* **the** Minkowski inequality

Fefferman and Stein's famous theorem,

*more usual:* the famous Fefferman–Stein theorem

7. In some expressions describing a noun, especially after “with” and “of”:

an algebra **with** unit  $e$ ; an operator **with** domain  $H^2$ ; a solution **with** vanishing Cauchy data; a cube **with** sides parallel to the axes; a domain **with** smooth boundary; an equation **with** constant coefficients; a function **with** compact support; random variables **with** zero expectation

the equation **of** motion; the velocity **of** propagation;

an element **of** finite order; a solution **of** polynomial growth;

a ball **of** radius 1; a function **of** norm  $p$

*But:* elements of **the** form  $f = \dots$

a Banach space with a weak symplectic form  $w$

two random variables with a common distribution

8. After forms of “have”:

It has  $\left\{ \begin{array}{l} \text{finite norm.} \\ \text{compact support.} \end{array} \right.$  *But:* It has  $\left\{ \begin{array}{l} \text{a finite norm not exceeding 1.} \\ \text{a compact support contained in } I. \end{array} \right.$

It has  $\left\{ \begin{array}{l} \text{rank } 2. \\ \text{cardinality } c. \\ \text{absolute value } 1. \\ \text{determinant zero.} \end{array} \right.$

*But:* It has  $\left\{ \begin{array}{l} \text{a zero of order at least } 2 \\ \text{at the origin.} \\ \text{a density } g. \\ \text{[Unless } g \text{ has appeared} \\ \text{earlier; then: It has density } g.] \end{array} \right.$

9. In front of the name of a mathematical discipline:

This idea comes from game theory (homological algebra).

*But:* in **the theory of** distributions

10. Other examples:

We can assume that  $G$  is **in diagonal form**.

Then  $A$  is deformed into  $B$  by pushing it **at constant speed** along the integral curves of  $X$ .

$G$  is now viewed as a set, **without group structure**.

## INFINITIVE

1. Indicating aim or intention:

**To prove** the theorem, we first let .....

We now apply (5)  $\left\{ \begin{array}{l} \text{to study the group of .....} \\ \text{to derive the following theorem.} \\ \text{to obtain an } x \text{ with norm not exceeding } 1. \end{array} \right.$

Here are some examples **to show** how .....

2. In constructions with “too” and “enough”:

This method is **too** complicated **to** be used here.

This case is important **enough to** be stated separately.

3. Indicating that one action leads to another:

We now apply Theorem 7 **to get**  $Nf = 0$ . [= ..... and we get  $Nf = 0$ ]

Insert (2) into (3) **to find** that .....

4. In constructions like “we may assume  $M$  to be .....”:

We may **assume**  $M$  **to be** compact.

We **define**  $K$  **to be** the section of  $H$  over  $S$ .

If we **take** the contour  $G$  **to lie** in  $U$ , then .....

We **extend**  $f$  **to be** homogeneous of degree 1.

The class  $A$  is defined by **requiring** all the functions  $f$  **to satisfy** .....

Partially order  $P$  by **declaring**  $X < Y$  **to mean** that .....

5. In constructions like “ $M$  is assumed to be .....”:

The map $M$	is <b>assumed</b> ⟨expected/found/considered/taken/ claimed⟩ <b>to be</b> open. will be <b>chosen to satisfy</b> (2). can be <b>taken to be</b> constant. can easily be <b>shown to have</b> ..... is also <b>found to be</b> of class $S$ .
-------------	---

This investigation is **likely to produce** good results.

[= It is very probable it will]

The close agreement of the six elements is **unlikely to be**  
a coincidence. [= probably not]

6. In the structure “for this to happen”:

**For this to happen**,  $F$  must be compact.

[= In order that this happens]

**For the last estimate to hold**, it is enough to assume .....

Then **for such a map to exist**, we must have .....

7. As the subject of a sentence:

**To see** that this is not a symbol is fairly easy.

[*Or*: It is fairly easy to see that .....

**To choose** a point at random in the interval  $[0, 1]$  is a conceptual  
experiment with an obvious intuitive meaning.

**To say** that  $u$  is maximal **means** simply that .....

*After expressions with “it”:*

**It is necessary** ⟨useful/very important⟩ **to consider** .....

**It makes sense to speak** of .....

**It is** therefore **of interest to look** at .....

8. After forms of “be”:

Our goal ⟨method/approach/procedure/objective/aim⟩ **is to find** .....

The problem ⟨difficulty⟩ here **is to construct** .....

9. With nouns and with superlatives, in the place of a relative clause:

The theorem **to be proved** is the following. [= which will be proved]

This will be proved by the method **to be described** in Section 6.

For other reasons, **to be discussed** in Chapter 4, we have to .....

He was **the first to propose** a complete theory of .....

They appear to be **the first to have suggested** the now accepted  
interpretation of .....

10. After certain verbs:

These properties **led him to suggest** that .....

Lax **claims to have obtained** a formula for .....

This map **turns out to satisfy** .....

At first glance  $M$  **appears to differ** from  $N$  in two major ways: .....

A more sophisticated argument **enables** one **to prove** that .....

[*Note*: “enable” requires “one”, “us” etc.]

He **proposed to study** that problem. [*Or*: He proposed studying .....

We **make**  $G$  **act** trivially on  $V$ .

**Let**  $f$  **satisfy** (2). [*Not*: “Let  $f$  satisfies”]

We **need to consider** the following three cases.

We **need not consider** this case separately.

[“need to” in affirmative clauses, without “to” in negative clauses; also note: “we only need to consider”, but: “we need only consider”]

## ING-FORM

1. As the subject of a sentence (note the absence of “the”):

**Repeating** the previous argument and **using** (3) **leads to** .....

Since **taking** symbols **commutes** with lifting,  $A$  is .....

**Combining** Proposition 5 and Theorem 7 **gives** .....

2. After prepositions:

**After making** a linear transformation, we may assume that .....

**In passing** from (2) to (3) we have ignored the factor  $n$ .

**In deriving** (4) we have made use of .....

**On substituting** (2) into (3) we obtain .....

**Before making** some other estimates, we prove .....

The trajectory  $Z$  enters  $X$  **without meeting**  $x = 0$ .

**Instead of using** the Fourier method we can multiply .....

**In addition to illustrating** how our formulas work, it provides .....

**Besides being** very involved, this proof gives no information on .....

This set is obtained **by letting**  $n \rightarrow \infty$ .

It is important to pay attention to domains of definition

**when trying to** .....

The following theorem is the key **to constructing** .....

The reason **for preferring** (1) to (2) is simply that .....

3. In certain expressions with “of”:

The **idea of combining** (2) and (3) came from .....

The **problem** considered there was that **of determining**  $WF(u)$  for .....

We use the **technique of extending** .....

This method has the **disadvantage of**

	<b>being</b> very involved.
	<b>requiring</b> that $f$ be positive.
	[ <i>Note</i> the infinitive.]

Actually,  $S$  has the much stronger **property of being** convex.

4. After certain verbs, especially with prepositions:

We **begin by analyzing** (3).

We **succeeded** (were successful) **in proving** (4).

[*Not*: “succeeded to prove”]

We next **turn to estimating** .....

They **persisted in investigating** the case .....

We are **interested in finding** a solution of .....

We were **surprised at finding out** that .....

[*Or*: surprised to find out]

Their study **resulted in proving** the conjecture for .....

The success of our method will **depend on proving** that .....

To compute the norm of ..... **amounts to finding** .....

We should **avoid using** (2) here, since .....

[*Not*: “avoid to use”]

We **put off discussing** this problem to Section 5.

It is **worth noting** that ..... [*Not*: “worth to note”]

It is worth while discussing here this phenomenon.

[*Or*: worth while to discuss; “worth while” with ing-forms is best avoided as it often leads to errors.]

It is an idea **worth carrying out**.

[*Not*: “worth carrying out”, *nor*: “worth to carry out”]

After **having finished proving** (2), we will turn to .....

[*Not*: “finished to prove”]

However, (2) **needs handling** with greater care.

One more case **merits mentioning** here.

In [7] he **mentions having proved** this for  $f$  not in  $S$ .

5. Present Participle in a separate clause (note that the subjects of the main clause and the subordinate clause must be the same):

We show that  $f$  satisfies (2), thus **completing** the analogy with .....

**Restricting** this to  $R$ , we can define .....

[*Not*: “Restricting ....., the lemma follows”. The lemma does not restrict!]

The set  $A$ , **being** the union of two intersecting continua, is connected.

6. Present Participle describing a noun:

We need only consider paths **starting** at 0.

We interpret  $f$  as a function with image **having** support in .....

We regard  $f$  as **being** defined on .....

7. In expressions which can be rephrased using “where” or “since”:

Now  $J$  is defined to equal  $Af$ , the function  $f$  **being** as in (3).

[= where  $f$  is .....

This is a special case of (4), the space  $X$  here **being**  $B(K)$ .

We construct three maps of the form (5), each of them **satisfying** (8).

Then  $\lim_t a(x, t) < 1$ , the limit **being assumed** to exist for every  $x$ .



The ideal is defined by  $m = \dots$ , it **being understood** that .....  
 Now,  $F$  **being** convex, we can assume that ..... [= since  $F$  is .....]  
 Hence  $F = \emptyset$  (it **being** impossible to make  $A$  and  $B$  intersect).  
 [= since it is impossible]

[Do not write “a function being an element of  $X$ ” if you mean  
 “a function which is an element of  $X$ ”.]

8. In expressions which can be rephrased as “the fact that  $X$  is .....”:

Note that  $M$  **being** cyclic implies  $F$  is cyclic.

The probability of  $X$  **being** rational equals  $1/2$ .

In addition to  $f$  **being** convex, we require that .....

## PASSIVE VOICE

1. Usual passive voice:

This theorem was proved by Milnor in 1976.

In items 2–6, passive voice structures replace sentences with subject “we” or impersonal constructions of other languages.

2. Replacing the structure “we do something”:

This identity **is established** by observing that .....

This difficulty **is avoided** above.

When this **is substituted** in (3), an analogous description of  $K$   
 is obtained.

Nothing **is assumed** concerning the expectation of  $X$ .

3. Replacing the structure “we prove that  $X$  is”:

The function  $M$  **is easily shown to have** .....  
**may be said to be** regular if .....

This equation **is known to hold** for .....

4. Replacing the construction “we give an object  $X$  a structure  $Y$ ”:

Note that  $E$  **can be given** a complex structure by .....

The letter  $A$  **is here given a bar** to indicate that .....

5. Replacing the structure “we act on something”:

This order behaves well when  $g$  **is acted upon** by an operator.

Hence  $F$  **can be thought of** as .....

So all the terms of (5) **are accounted for**.

The preceding observation, when **looked at** from a more general  
 point of view, leads to .....

In the physical context already **referred to**,  $K$  is .....

6. Meaning “which will be (proved etc.)”:

Before stating the result **to be proved**, we give .....

This is a special case of convolutions **to be introduced** in Chapter 8.

We conclude with two simple lemmas **to be used** mainly in .....

**QUANTIFIERS**

This implies that  $A$  contains  $\left\{ \begin{array}{l} \text{all open subsets of } U. \\ \text{all } y \text{ with } Gy = 1. \end{array} \right.$

Let  $B$  be the collection of  $\left\{ \begin{array}{l} \text{all transforms } F \text{ of the form .....} \\ \text{all } A \text{ such that .....} \end{array} \right.$

In this way  $F$  is defined at **all** points of  $X$ .

This holds for **all**  $n \neq 0$  (for **all**  $m$  which have ...../for **all** other  $m$ / for **all but** a finite number of indices  $i$ )

The domain  $X$  contains **all the** boundary except the origin.

The integral is taken over **all of**  $X$ .

Hence  $E, F$  and  $G$   $\left\{ \begin{array}{l} \text{all extend to a neighbourhood of } U. \\ \text{all have their supports in } U. \\ \text{are all zero at } x. \\ \text{are all equal.} \end{array} \right.$

There exist functions  $R$ , **all of whose** poles are in  $U$ , with .....

Each of the following nine conditions implies **all the others**.

Such an  $x$  exists iff **all the** intervals  $A_x$  have .....

For **every**  $g$  in  $X$  (not in  $X$ ) there exists an  $N$  .....

[But: for all  $f$  and  $g$ , for any two maps  $f$  and  $g$ ;  
“every” is followed by a *singular* noun.]

To **every**  $f$  there corresponds a unique  $g$  such that .....

**Every** invariant subspace of  $X$  is of the form .....

[Do not write: “Every subspace is not of the form .....”  
if you mean: “No subspace is of the form .....”;  
“every” must be followed by an *affirmative*  
statement.]

Thus  $f \neq 0$  at **almost every** point of  $X$ .

Since  $A_n = 0$  for **each**  $n$ , .....

[Each = every, considered separately]

**Each** term in this series is either 0 or 1.

Consequently,  $F$  is bounded on **each** bounded set.

**Each** of these four integrals is finite.

These curves arise from ....., and **each** consists of .....

There remain four intervals of length  $1/16$  **each**.

Thus  $X$  assumes values  $0, 1, \dots, 9$ , **each** with probability  $1/10$ .

The functions  $F_1, \dots, F_n$  are **each** defined in the interval  $[0, 1]$ .

Those  $n$  disjoint boxes are translates **of each other**.

If  $K$  is now **any** compact subset of  $H$ , there exists .....

[Any = whatever you like; write “for all  $x$ ”, “for every  $x$ ” if you just mean a quantifier.]

Every measure can be completed, so whenever it is convenient, we may assume that **any** given measure is complete.

**There is** a subsequence such that .....

**There exists** an  $x$  with .....

[Or: there exists  $x$ , but: there is an  $x$ ]

**There are** sets satisfying (2) but not (3).

**There is** only one such  $f$ .

**There is** a unique function  $f$  such that .....

Each  $f$  lies in  $zA$  for **some**  $A$  (at least one  $A$ /  
exactly one  $A$ /at most one  $A$ ).

Note that **some of** the  $X_n$  may be repeated.

Thus  $F$  has **no** pole in  $U$  (hence **none** in  $K$ ). [Or: no poles]

Call a set dense if its complement contains **no** nonempty open subset.

If **no two** members of  $A$  have an element in common, then .....

**No two** of the spaces  $X$ ,  $Y$ , and  $Z$  are isomorphic.

It can be seen that **no**  $x$  has more than one inverse.

In other words, for **no** real  $x$  does  $\lim F_n(x)$  exist.

[Note the inversion after the negative clause.]

If there is **no** bounded functional such that .....

..... provided **none of** the sums is of the form .....

Let  $A_n$  be a sequence of positive integers **none of which** is 1 less than a power of two.

If there is an  $f$  such that ....., set ....., If **there are** (is) **none**, define .....

**None of these** are (is) possible.

**Both**  $f$  and  $g$  are obtained by .....

[Or:  $f$  and  $g$  are both obtained]

For **both**  $C^\infty$  and analytical categories, .....

It behaves covariantly with respect to maps of **both**  $X$  and  $G$ .

We now apply (3) to **both** sides of (4).

**Both** (these/the) conditions are restrictions only on .....

[Note: “the” and “these” after “both”]

It lies on no segment **both of whose** endpoints are in  $K$ .

Two consecutive elements do not belong **both to**  $A$   
or **both to**  $B$ .

**Both its** sides are convex. [Or: Its sides are both convex.]

Let  $B$  and  $C$  be nonnegative numbers, not **both** 0.

Choose points  $x$  in  $M$  and  $y$  in  $N$ , **both** close to  $z$ , and .....

We show how this method works in two cases.

In **both** (In each),  $C$  is .....

In **either** case, it is clear that ..... [= In both cases]

Each  $f$  can be expressed in **either of** the forms (1) and (2).

[= in any of the two forms]

The density of  $X + Y$  is given by **either of** the two integrals.

The two classes coincide if  $X$  is compact. In that case we write  $C(X)$  for **either of** them.

**Either**  $f$  or  $g$  must be bounded.

Let  $u$  and  $v$  be two distributions **neither of** which is .....

[Use “neither” when there are *two* alternatives.]

This is true for **neither of** the two functions.

**Neither** statement is true.

In **neither** case can  $f$  be smooth.

[Note the inversion after the negative clause.]

He proposes two conditions, but **neither** is satisfactory.

## NUMBER, QUANTITY, SIZE

### 1. Cardinal numbers:

Hence  $A$  and  $B$  are also  $F$ -functions, any **two** of  $A$ ,  $B$ , and  $C$  being independent.

the multi-index with  $\left\{ \begin{array}{l} \text{all entries } \mathbf{zero} \text{ except the } k\text{th which is } \mathbf{one} \\ \text{the last } k \text{ entries } \mathbf{zero} \end{array} \right.$

This shows that there are no **two** points  $a$  and  $b$  such that .....

There are **three** that the reader must remember. [= three of them]

We have defined  $A$ ,  $B$ , and  $C$ , and **the three** sets satisfy .....

For **the two** maps defined in Section 3, .....

[“The” if only two maps are defined there.]

Clearly,  $R$  is concentrated at **the**  $n$  points  $x_1, \dots, x_n$  defined above.

for **at least**  $\langle$ **at most** $\rangle$  one  $k$ ; with norm **at least equal to** 2

There are **at most 2 such**  $r$  in  $(0, 1)$ .

There is **a unique** map satisfying (4).

Equation (4) has **a unique** solution  $g$  for each  $f$ .

*But:* it has **the unique** solution  $g = ABf$ .

Problem (4) has **one and only one** solution.

**Precisely**  $r$  of the intervals are closed.

In Example 3 only **one of** the  $x_j$  is positive.

If  $p = 0$  then there are **an additional**  $m$  arcs.

### 2. Ordinal numbers:

**The first two** are simpler than **the third**.

Let  $S_i$  be **the first of** the remaining  $S_j$ .

**The  $n$ th** trial is the last.

It follows that  $X_1$  appears at **the**  $(k + 1)$ **th** place.

The gain **up to and including the  $n$ th** trial is .....

The elements of **the third and fourth** rows are in  $I$ .  
 [Note the plural.]

Therefore  $F$  has a zero of **at least third** order at  $x$ .

3. Fractions:

**Two-thirds** of its diameter is covered by .....

*But:* **Two-thirds** of the gamblers are ruined.

Obviously,  $G$  is **half** the sum of the positive roots.  
 [Note: Only “half” can be used with or without “of”.]

On the average, about **half** the list will be tested.  
 But  $J$  contains an interval of **half** its length in which .....

Note that  $F$  is greater by **a half** (a third).  
 The other player is half (one third) as fast.  
 We divide  $J$  **in half**.  
 All sides were increased by the same **proportion**.  
 About **40 percent** of the energy is dissipated.  
 A positive **percentage** of summands occurs in all  $k$  partitions.

4. Smaller (greater) than:

Observe that $n$ is	<b>greater</b> (less) <b>than</b> $k$ . <b>much</b> (substantially) <b>greater</b> than $k$ . <b>no greater</b> (smaller) <b>than</b> $k$ . <b>greater</b> (less) <b>than or equal to</b> $k$ . [Not: “greater or equal to”] <b>strictly less than</b> $k$ .
---------------------	---

All points at a distance **less than**  $K$  from  $A$  satisfy (2).  
 We thus obtain a graph of **no more than**  $k$  edges.

This set has **fewer** elements than  $K$  has.  
**no fewer than** twenty elements.

Therefore  $F$  can have no jumps **exceeding**  $1/4$ .  
 The degree of  $P$  **exceeds** that of  $Q$ .  
 Find the density of **the smaller of**  $X$  and  $Y$ .  
**The smaller of the two** satisfies .....

It is dominated (bounded/estimated/majorized) by .....

5. How much smaller (greater):

25 is **3 greater** than 22; 22 is **3 less** than 25.  
 Let  $a_n$  be a sequence of positive integers none of which is **1 less** than a power of two.  
 The degree of  $P$  **exceeds** that of  $Q$  by **at least 2**.  
 Consequently,  $f$  is **greater by a half** (a third).  
 It follows that  $C$  is **less than a third** of the distance between .....

Within  $I$ , the function  $f$  **varies** (oscillates)  
by **less than**  $l$ .

The upper and lower limits of  $f$  **differ by at most** 1.

We thus have in  $A$  **one** element **too many**.

On applying this argument  $k$  **more times**, we obtain .....

This method is recently **less and less** used.

A succession of **more and more** refined discrete models.

6. How many times as great:

twice (ten times/one third) **as long as**; half as big as

The longest edge is at most 10 times as long as the shortest one.

Now  $A$  has **twice as many** elements as  $B$  has.

Clearly,  $J$  contains a subinterval **of half its length** in which .....

Observe that  $A$  has four times the radius of  $B$ .

The diameter of  $L$  is  $1/k$  times (twice) **that of**  $M$ .

7. Multiples:

The  $k$ -fold integration by parts shows that .....

We have shown that  $F$  covers  $M$  **twofold**.

It is bounded by **a multiple of**  $t$  (a constant times  $t$ ).

This distance is less than **a constant multiple of**  $d$ .

Note that  $G$  acts on  $H$  as **a multiple**, say  $n$ , of  $V$ .

8. Most, least, greatest, smallest:

Evidently,  $F$  has **the most** (the fewest) points when .....

In **most** cases it turns out that .....

**Most of** the theorems presented here are original.

The proofs are, **for the most part**, only sketched.

**Most probably**, his method will prove useful in .....

What **most** interests us is whether .....

The **least** such constant is called the norm of  $f$ .

This is **the least** useful of the four theorems.

The method described above seems to be **the least** complex.

That is **the least** one can expect.

The elements of  $A$  are comparatively big, but **least** in number.

None of those proofs is easy, and John's **least of all**.

The best estimator is a linear combination  $U$  such that  
 $\text{var } U$  is (the) **smallest possible**.

The expected waiting time is **smallest** if .....

Let  $L$  be **the smallest number** such that .....

Now,  $F$  has **the smallest** norm among all  $f$  such that .....

It is **the largest of** the functions which occur in (3).

There exists **a smallest** algebra with this property.

Find the **second largest** element in the list  $L$ .

9. Many, few, a number of:

There are [Note the plural.]	a <b>large number</b> of illustrations. only a <b>finite number</b> of $f$ with $Lf = 1$ . a <b>small number</b> of exceptions. an <b>infinite number</b> of sets .....
------------------------------------	--

Ind  $c$  is **the number of times** that  $c$  winds around 0.

We give **a number of** results concerning ..... [= some]

This may happen in **a number of** cases.

They correspond to the values of **a countable number of** invariants.

..... for **all  $n$  except a finite number** (for **all but finitely many  $n$** ).

Thus  $Q$  contains **all but a countable number** of the  $f^i$ .

There are only **countably many** elements  $q$  of  $Q$  with  $\text{dom } q = S$ .

The theorem is fairly general. There are, however, **numerous** exceptions.

**A variety** of other characteristic functions can be constructed in this way.

There are **few** exceptions to this rule. [= not many]

**Few** of various existing proofs are constructive.

He accounts for all the major achievements in topology over **the last few** years.

The generally accepted point of view in this domain of science seems to be changing **every few** years.

There are **a few** exceptions to this rule. [= some]

Many interesting examples are known. We now describe **a few of** these.

Only **a few of** those results have been published before.

**Quite a few of** them are now widely used.

[= A considerable number]

10. Equality, difference:

$A$  equals  $B$  or  $A$  is equal to  $B$  [Not: " $A$  is equal  $B$ "]

The Laplacian of  $g$  is  $4r > 0$ .

Then  $r$  is about  $kn$ .

The inverse of  $FG$  is  $GF$ .

The norms of  $f$  and  $g$  coincide.

Therefore  $F$  has the same number of zeros and poles in  $U$ .

They **differ by** a linear term (by a scale factor).

The differential of  $f$  is **different from** 0.

Each member of  $G$  **other than**  $g$  is .....

Lemma 2 shows that  $F$  is not identically 0.

Let  $a$ ,  $b$  and  $c$  be **distinct** complex numbers.

Each  $w$  is  $Pz$  for precisely  $m$  distinct values of  $z$ .

Functions which are equal a.e. are indistinguishable as far as integration is concerned.

## 11. Numbering:

Exercises 2 to 5 furnish other applications of this technique.

[*Amer.*: Exercises 2 through 5]

in the third and fourth rows

the derivatives up to order  $k$

from row  $k$  onwards

the odd-numbered terms

in lines 16–19

the next-to-last column

the last paragraph but one of the previous proof

The matrix with  $\left| \begin{array}{l} 1 \text{ in the } (i, j) \text{ entry and zero elsewhere} \\ \text{all entries zero except for } N - j \text{ at } (N, j) \end{array} \right.$

This is  $\left| \begin{array}{l} \text{hinted at in Sections 1 and 2.} \\ \text{quoted on page 36 of [4].} \end{array} \right.$

## HOW TO AVOID REPETITION

### 1. Repetition of nouns:

Note that the continuity of  $f$  implies **that** of  $g$ .

The passage from Riemann's theory to **that** of Lebesgue is .....

The diameter of  $F$  is about twice **that** of  $G$ .

His method is similar to **that** used in our previous paper.

The nature of this singularity is the same as **that**

which  $f$  has at  $x = 0$ .

Our results do not follow from **those** obtained by Lax.

One can check that the metric on  $T$  is **the one** we have just described.

It follows that  $S$  is the union of two disks. Let  $D$  be **the one** that contains .....

The cases  $p = 1$  and  $p = 2$  will be **the ones** of interest to us.

We prove a uniqueness result, similar to **those** of the preceding section.

Each of the functions on the right of (2) is **one**  
to which .....

Now,  $F$  has many points of continuity. Suppose  $x$  is **one**.

In addition to a contribution to  $W_1$ , there may be **one**  
to  $W_2$ .

We now prove that the constant  $pq$  cannot be replaced by  
a smaller **one**.

Consider the differences between these integrals and

**the corresponding ones** with  $f$  in place of  $g$ .

The geodesics (4) are **the only ones** that realize the distance between  
their endpoints.

On account of the estimate (2) and similar **ones** which can be .....



We may replace  $A$  and  $B$  by whichever is the larger of **the two**.  
[*Not*: “the two ones”]

This inequality applies to conditional expectations as well as to ordinary **ones**.

One has to examine the equations (4). If **these** have no solutions, then .....

Thus  $D$  yields operators  $D^+$  and  $D^-$ . **These** are formal adjoints of each other.

This gives rise to the maps  $F_i$ . All the other maps are suspensions of **these**.

So  $F$  is the sum of  $A$ ,  $B$ ,  $C$  and  $D$ . The last two of **these** are zero.

Both  $f$  and  $g$  are connected, but **the latter** is in addition compact.  
[The latter = the second of *two* objects]

Both  $AF$  and  $BF$  were first considered by Banach, but only **the former** is referred to as the Banach map, **the latter** being called the Hausdorff map.

We have thus proved Theorems 1 and 2, **the latter** without using .....

Since the vectors  $G_i$  are orthogonal to **this last** space, .....

As a consequence of **this last** result, .....

Let us consider sets of the type (1), (2), (3) and (4).

**These last two** are called .....

We shall now describe a general situation in which **the last-mentioned** functionals occur naturally.

## 2. Repetition of adjectives, adverbs or phrases like “ $x$ is .....”:

If  $f$  and  $g$  are measurable functions, then **so are**  $f + g$  and  $f \cdot g$ .

The union of measurable sets is a measurable set; **so is** the complement of every measurable set.

The group  $G$  is compact and **so is** its image under  $f$ .

It is of the same fundamental importance in analysis **as is** the construction of .....

Note that  $F$  is bounded but **is not** necessarily **so** after division by  $G$ .

Show that there are many **such**  $Y$ .

There is only one **such** series for each  $y$ .

**Such an**  $h$  is obtained by .....

## 3. Repetition of verbs:

A geodesic which meets  $bM$  **does so** either transversally or .....

This will hold for  $x > 0$  if it **does** for  $x = 0$ .

Note that we have not required that ....., and we shall not **do so** except when explicitly stated.

The integral might not converge, but it **does so** after .....

We will show below that the wave equation can be put in this form,  
**as can** many other systems of equations.

The elements of  $L$  are not in  $S$ , **as they are** in the proof of .....

#### 4. Repetition of whole sentences:

**The same is true** for  $f$  in place of  $g$ .

**The same being true** for  $f$ , we can .....

**The same holds** for  $\langle$ applies to $\rangle$  the adjoint map.

We shall assume that **this is the case**.

**Such was the case** in (2).

The  $L^2$  theory has more symmetry **than is the case**  
in  $L^1$ .

Then either ..... or ..... **In the latter**  $\langle$ former $\rangle$  **case**, .....

For  $k$  **this is no longer true**.

**This is not true of** (2).

**This is not so** in other queuing processes.

If **this is so**, we may add .....

If  $f_i \in L$  and if  $F = f_1 + \dots + f_n$  then  $F \in H$ , and every  
 $F$  is **so** obtained.

We would like to .....

If  $U$  is open, **this** can be done.

On  $S$ , **this** gives the ordinary topology of the plane.

Note that **this** is not equivalent to .....

[Note the difference between “this” and “it”: you say “*it* is not  
equivalent to” if you are referring to some object explicitly  
mentioned in the preceding sentence.]

Consequently,  $F$  **has the stated**  $\langle$ desired/claimed $\rangle$  **properties**.

## WORD ORDER

*General remarks:* The normal order is: subject + verb + direct object + adverbs in  
the order manner-place-time.

Adverbial clauses can also be placed at the beginning of a sentence, and some adverbs  
always come between subject and verb. Subject almost always precedes verb,  
except in questions and some negative clauses.

### 1. ADVERBS

1a. Between subject and verb, but after forms of “be”; in compound tenses  
after first auxiliary

#### • *Frequency adverbs:*

This has **already** been proved in Section 8.

This result will **now** be derived computationally.

Every measurable subset of  $X$  is **again** a measure space.

We **first** prove a reduced form of the theorem.

There has **since** been little systematic work on .....  
It has **recently** been pointed out by Fix that .....  
It is **sometimes** difficult to .....  
This **usually** implies further conclusions about  $f$ .  
It **often** does not matter whether .....

- *Adverbs like “also”, “therefore”, “thus”:*

Our presentation is **therefore** organized in such a way that .....  
The sum in (2), though formally infinite, is **therefore** actually finite.  
One must **therefore** also introduce the class of .....  
But  $C$  is connected and is **therefore** not the union of .....

These properties, with the exception of (1), **also** hold  
for  $t$ .

We will **also** leave to the reader the verification that .....  
It will **thus** be sufficient to prove that .....  
So (2) implies (3), since one would **otherwise** obtain .....

The order of several topics has **accordingly** been changed.

- *Emphatic adverbs (clearly, obviously, etc.):*

It would **clearly** have been sufficient to assume that .....  
But  $F$  is **clearly** not an  $I$ -set.  
Its restriction to  $N$  is **obviously** just  $f$ .  
This case must **of course** be excluded.  
The theorem **evidently** also holds if  $x = 0$ .

The crucial assumption is that the past history  
**in no way** influences .....

We did not **really** have to use the existence of  $T$ .  
The problem is to decide whether (2) **really** follows  
from (1).

The proof is now **easily** completed.

The maximum is **actually** attained at some point of  $M$ .

We then **actually** have ..... [= We have even more]

At present we will **merely** show that .....

A stronger result is **in fact** true.

Throughout integration theory, one **inevitably** encounters  $\infty$ .

But  $H$  itself can **equally well** be a member of  $S$ .

- 1b. After verb—most adverbs of manner:

We conclude **similarly** that .....

One sees **immediately** that .....

Much relevant information can be obtained **directly** from (3).

This difficulty disappears **entirely** if .....

This method was used **implicitly** in random walks.

1c. After an object if it is short:

We will prove the theorem **directly** without using the lemma.

*But:* We will prove **directly** a theorem stating that .....

This is true for every sequence that shrinks to  $x$  **nicely**.

Define  $Fg$  **analogously** as the limit of .....

Formula (2) defines  $g$  **unambiguously** for every  $g'$ .

1d. At the beginning—adverbs referring to the whole sentence:

**Incidentally**, we have now constructed .....

**Actually**, Theorem 3 gives more, namely .....

**Finally**, (2) shows that  $f = g$ . [*Not:* “At last”]

**Nevertheless**, it turns out that .....

**Next**, let  $V$  be the vector space of .....

**More precisely**,  $Q$  consists of .....

**Explicitly** (**Intuitively**), this means that .....

**Needless to say**, the boundedness of  $f$  was assumed only for simplicity.

**Accordingly**, either  $f$  is asymptotically dense or .....

1e. In front of adjectives—adverbs describing them:

a **slowly varying** function

**probabilistically significant** problems

a method **better suited** for dealing with .....

The maps  $F$  and  $G$  are **similarly obtained** from  $H$ .

The function  $F$  has a **rectangularly shaped** graph.

Three-quarters of this area is covered by **subsequently chosen** cubes. [Note the singular.]

1f. “only”

We need the openness **only** to prove the following.

It reduces to the statement that **only** for the distribution  $F$  do the maps  $F_i$  satisfy (2). [Note the inversion.]

In this chapter we will be concerned **only** with .....

In (3) the  $X_j$  assume the values 0 and 1 **only**.

If (iii) is required for finite unions **only**, then .....

We need **only** require (5) to hold for bounded sets.

The proof of (2) is similar, and will **only** be indicated briefly.

To prove (3), it **only** remains to verify .....

## 2. ADVERBIAL CLAUSES

2a. At the beginning:

**In testing** the character of ....., it is sometimes difficult to .....

**For**  $n = 1, 2, \dots$ , consider a family of .....

2b. At the end (normal position):

The averages of  $F_n$  become small **in small neighbourhoods** of  $x$ .

2c. Between subject and verb, but after first auxiliary—only short clauses:

The observed values of  $X$  will **on average** cluster around .....

This could **in principle** imply an advantage.

For simplicity, we will **for the time being** accept as  $F$  only  $C^2$  maps.

Accordingly we are **in effect** dealing with .....

The knowledge of  $f$  is **at best** equivalent to .....

The stronger result is **in fact** true.

It is **in all respects** similar to matrix multiplication.

2d. Between verb and object if the latter is long:

It suffices **for our purposes** to assume .....

To a given density on the line there corresponds **on the circle**  
the density given by .....

### 3. INVERSION AND OTHER PECULIARITIES

3a. Adjective or past participle after a noun:

Let  $Y$  be the complex  $X$  with the origin **removed**.

Theorems 1 and 2 **combined** give a theorem .....

We now show that  $G$  is in the symbol class **indicated**.

We conclude by the part of the theorem **already proved** that .....

The bilinear form **so defined** extends to .....

Then for  $A$  **sufficiently small** we have .....

By queue length we mean the number of customers **present**  
including the customer **being served**.

The description is the same with the roles of  $A$  and  $B$  **reversed**.

3b. Direct object or adjectival clause placed farther than usual—when they are long:

We must **add** to the right-hand side of (3) **the probability** that .....

This is equivalent to **defining** in the  $z$ -plane **a density** with .....

Let  $F$  be the **restriction** to  $D$  **of** the unique linear map .....

The **probability** at birth **of** a lifetime exceeding  $t$  is at most .....

3c. Inversion in some negative clauses:

We do not assume that ....., **nor do** we assume a priori that .....

**Neither is** the problem simplified by assuming  $f = g$ .

The “if” part now follows from (3), since **at no point can**  $S$  exceed  
the larger of  $X$  and  $Y$ .

The fact that **for no  $x$  does**  $Fx$  contain  $y$  implies that .....

**In no case does** the absence of a reference imply any claim to  
originality on my part.

3d. Inversion—other examples:

But  $F$  is compact and **so is**  $G$ .

If  $f, g$  are measurable, then **so are**  $f + g$  and  $f \cdot g$ .

**Only** for  $f = 1$  | **can** one expect to obtain .....  
| **does** that limit exist.

3e. Adjective in front of forms of “be”—for emphasis:

By far **the most important** is the case where .....

**Much more subtle** are the following results of John.

**Essential** to the proof are certain topological properties of  $M$ .

3f. Subject coming sooner than in some other languages:

**Equality** occurs in (1) iff  $f$  is constant.

The natural **question** arises whether it is possible to .....

In the following applications **use** will be made of .....

Recently **proofs** have been constructed which use .....

3g. Incomplete clause at the beginning or end of a sentence:

**Put differently**, the moments of arrival of the lucky customers constitute a renewal process.

**Rather than discuss** this in full generality, let us look at .....

It is important that the tails of  $F$  and  $G$  are of comparable magnitude, **a statement** made more precise by the following inequalities.

## WHERE TO INSERT A COMMA

*General rules:* Do not over-use commas—English usage requires them less often than in many other languages. Do not use commas around a clause that defines (limits, makes more precise) some part of a sentence. Put commas before and after non-defining clauses (i.e. ones which can be left out without damage to the sense). Put a comma where its lack may lead to ambiguity, e.g. between two symbols.

1. Comma not required:

We shall now prove that  $f$  is proper.

The fact that  $f$  has radial limits was proved in [4].

It is reasonable to ask whether this holds for  $g = 1$ .

Let  $M$  denote the set of all paths that satisfy (2).

There is a polynomial  $P$  such that  $Pf = g$ .

The element given by (3) is of the form (5).

Let  $M$  be the manifold to whose boundary  $f$  maps  $K$ .

Take an element all of whose powers are in  $S$ .

We call  $F$  proper if  $G$  is dense.

There exists a  $D$  such that  $D \sim H$  whenever  $H \sim G$ .

Therefore  $F(x) = G(x)$  for all  $x \in X$ .

Let  $F$  be a nontrivial continuous linear operator in  $V$ .

## 2. Comma required:

The proof of (3) depends on the notion of  $M$ -space, which has already been used in [4].

We will use the map  $H$ , which has all the properties required.

There is only one such  $f$ , and (4) defines a map from .....

In fact, we can do even better.

In this section, however, we will not use it explicitly.

Moreover,  $F$  is countably additive.

Finally, (d) and (e) are consequences of (4).

Nevertheless, he succeeded in proving that .....

Conversely, suppose that .....

Consequently, (2) takes the form .....

In particular,  $f$  also satisfies (1).

Guidance is also given, whenever necessary or helpful, on further reading.

This observation, when looked at from a more general point of view, leads to .....

It follows that  $f$ , being convex, cannot satisfy (3).

If  $e = 1$ , which we may assume, then .....

We can assume, by decreasing  $k$  if necessary, that .....

Then (5) shows, by Fubini's theorem, that .....

Put this way, the question is not precise enough.

Being open,  $V$  is a union of disjoint boxes.

This is a special case of (4), the space  $X$  here being  $B(K)$ .

In [2],  $X$  is assumed to be compact.

For all  $x$ ,  $G(x)$  is convex.

[Comma between two symbols.]

In the context already referred to,  $K$  is the complex field.

[Comma to avoid ambiguity.]

## 3. Comma optional:

By Theorem 2, there exists an  $h$  such that .....

For  $z$  near 0, we have .....

If  $h$  is smooth, then  $M$  is compact.

Since  $h$  is smooth,  $M$  is compact.

It is possible to use (4) here, but it seems preferable to .....

This gives (3), because (since) we may assume .....

Integrating by parts, we obtain .....

The maps  $X$ ,  $Y$ , and  $Z$  are all compact.

We have  $X = FG$ , where  $F$  is defined by .....

Thus (Hence/Therefore), we have .....

## HYPHENATION

### 1. Non(-):

Write consistently either

nontrivial, nonempty, nondecreasing, nonnegative, or  
non-trivial, non-empty, non-decreasing, non-negative.

[*But*: non-locally convex, non-Euclidean]

### 2. Hyphen required:

one-parameter group

two-stage computation

$n$ -fold integration

out-degree

global-in-time solution [But: solution global in time]

### 3. Hyphen optional:

right hand side or right-hand side

second order equation or second-order equation

selfadjoint or self-adjoint

halfplane or half-plane

seminorm or semi-norm

a blow-up, a blow up, or a blowup [But: to blow up]

the  $n$ th element or the  $n$ -th element

## SOME TYPICAL ERRORS

### 1. Spelling errors:

Spelling should be either British or American throughout:

*Br.*: colour, neighbourhood, centre, fibre, labelled, modelling

*Amer.*: color, neighborhood, center, fiber, labeled, modeling

“an unified approach”  $\rightsquigarrow$  a unified approach

“a  $M$  such that”  $\rightsquigarrow$  an  $M$  such that

[Use  $a$  or  $an$  according to pronunciation.]

“preceeding”  $\rightsquigarrow$  preceding

“occuring”  $\rightsquigarrow$  occurring

“developped”  $\rightsquigarrow$  developed

“loosing”  $\rightsquigarrow$  losing

“it’s norm”  $\rightsquigarrow$  its norm

### 2. Grammatical errors:

“Let  $f$  denotes”  $\rightsquigarrow$  Let  $f$  denote

“Most of them is”  $\rightsquigarrow$  Most of them are

“There is a finite number of”  $\rightsquigarrow$  There are a finite number of



“In 1964 Lax has shown”  $\rightsquigarrow$  In 1964 Lax showed  
 [Use the past tense if a date is given.]

“the Taylor’s formula”  $\rightsquigarrow$  Taylor’s formula [Or: the Taylor formula]

“the section 1”  $\rightsquigarrow$  Section 1

“Such map exists”  $\rightsquigarrow$  Such a map exists [But: for every such map]

“in case of smooth norms”  $\rightsquigarrow$  in the case of smooth norms

“We are in the position to prove”  $\rightsquigarrow$  We are in a position to prove

“We now give few examples” [= not many]  
 $\rightsquigarrow$  We now give a few examples [= some]

“ $F$  is equal  $G$ ”  $\rightsquigarrow$   $F$  is equal to  $G$  [Or:  $F$  equals  $G$ ]

“ $F$  is greater or equal to  $G$ ”  $\rightsquigarrow$   $F$  is greater than or equal to  $G$

“This is precised by”  $\rightsquigarrow$  This is made more precise by

“This allows to prove”  $\rightsquigarrow$  This allows us to prove

“This makes clear that”  $\rightsquigarrow$  This makes it clear that

“The first two ones are”  $\rightsquigarrow$  The first two are

“a not dense set”  $\rightsquigarrow$  a non-dense set  
 [But: This set is not dense]

“Since  $f = 0$ , then  $M$  is closed”  
 $\rightsquigarrow$  Since  $f = 0$ , it follows that  $M$  is closed

“....., as it is shown in Sec. 2”  $\rightsquigarrow$  ....., as is shown in Sec. 2

“Every function being an element of  $X$  is convex”  
 $\rightsquigarrow$  Every function which is an element of  $X$  is convex

“Every  $f$  is not convex”  $\rightsquigarrow$  No  $f$  is convex

“Setting  $n = p$ , the equation can be solved by .....”  
 $\rightsquigarrow$  Setting  $n = p$ , we can solve the equation by .....”  
 [Because we set.]

“We have ⟨get/obtain⟩ that  $B$  is empty”  
 $\rightsquigarrow$  We see ⟨know/conclude/deduce/find/infer⟩ that  $B$  is empty

### 3. Wrong word used:

“Summing (2) and (3) by sides”  $\rightsquigarrow$  Summing (2) and (3)

“In the first paragraph”  $\rightsquigarrow$  In the first section

“which proves our thesis”  
 $\rightsquigarrow$  which proves our assertion ⟨conclusion/statement⟩  
 [thesis = dissertation]

“to this aim”  $\rightsquigarrow$  to this end

“At first, note that”  $\rightsquigarrow$  First, note that

“At last,  $C$  is dense because”  $\rightsquigarrow$  Finally,  $C$  is dense because

“for every two elements”  $\rightsquigarrow$  for any two elements

“....., what completes the proof”  $\rightsquigarrow$  ....., which completes the proof

“....., what is impossible”  $\rightsquigarrow$  ....., which is impossible

“We denote it shortly by  $c$ ”  $\rightsquigarrow$  We denote it briefly by  $c$

“This map verifies (2)”  $\rightsquigarrow$  This map satisfies (2)

“continuous in the point  $x$ ”  $\rightsquigarrow$  continuous at  $x$

“disjoint with  $B$ ”  $\rightsquigarrow$  disjoint from  $B$

“equivalent with  $B$ ”  $\rightsquigarrow$  equivalent to  $B$

“independent on  $B$ ”  $\rightsquigarrow$  independent of  $B$

[*But*: depending on  $B$ ,  
independence from  $B$ ]

“similar as  $B$ ”  $\rightsquigarrow$  similar to  $B$

similarly as in Sec. 2  $\rightsquigarrow$ 

similarly to Sec. 2
as (just as) in Sec. 2
as is the case in Sec. 2
in much the same way as
in Sec. 2

“on Fig. 3”  $\rightsquigarrow$  in Fig. 3

“in the end of Sec. 2”  $\rightsquigarrow$  at the end of Sec. 2

#### 4. Wrong word order:

“a bounded by 1 function”  $\rightsquigarrow$  a function bounded by 1

“the described above condition”  $\rightsquigarrow$  the condition described above

“the obtained solution”  $\rightsquigarrow$  the solution obtained

“the mentioned map”  $\rightsquigarrow$  the map mentioned

[*But*: the above-mentioned map]

“the both conditions”  $\rightsquigarrow$  both conditions, both the conditions

“its both sides”  $\rightsquigarrow$  both its sides

“the three first rows”  $\rightsquigarrow$  the first three rows

“the two following sets”  $\rightsquigarrow$  the following two sets

“This map we denote by  $f$ ”  $\rightsquigarrow$  We denote this map by  $f$

“Only for  $x = 1$  the limit exists”  $\rightsquigarrow$  Only for  $x = 1$  does the limit exist

“For no  $x$  the limit exists”  $\rightsquigarrow$  For no  $x$  does the limit exist

## INDEX

- a, an, 23, 46  
accordingly, 13  
actually, 19, 41  
adjectival clauses, 9  
adverbial clauses, 42  
adverbs, 40  
a few, 37, 47  
all, 32  
also, 41  
a number of, 37, 46  
any, 33  
as, 15, 18, 40  
as is, 39, 47  
at first, 47  
at last, 42, 47  
avoid, 30  
  
because, 15  
being, 9, 30, 47  
both, 33, 48  
brackets, 8  
briefly, 7, 48  
  
cardinal numbers, 34  
case, 40, 47  
contradiction, 14, 18  
  
denote, 7  
depending on, 8  
differ, 36, 37  
disjoint from, 48  
distinct, 37  
  
each, 32  
either, 34  
enable, 29  
enough, 8, 27  
equal, 37, 47  
every, 32, 47  
  
few, 37, 47  
fewer, 35  
finally, 42, 47  
finish, 30  
*k*-fold, 36  
following, 13, 19, 20, 47  
for, 11, 28  
former, 39, 40  
for short, 7  
fractions, 35  
  
generality, 10  
greater, 35  
  
half, 35  
have, 26  
“have that”, 15, 16, 47  
hence, 15, 45  
  
if necessary, 11  
imperative, 16  
in a position, 17, 47  
independent of, 8, 48  
induction, 14  
in fact, 13  
infinitive, 10, 17, 27, 29  
introduction, 4  
inversion, 9, 10, 33, 42, 43, 48  
it, 18, 19, 28, 40  
it follows that, 15, 47  
  
largest, 36  
last but one, 38  
latter, 39, 40  
least, 34, 36  
less, 35  
let, 46  
likely, 28  
  
matrices, 38  
more, 36  
most, 34, 36, 46  
multiple, 36  
  
need, 17, 29  
neither, 9, 34, 43  
next-to-last, 38  
no, 33, 48  
no greater, 35  
non(-), 46, 47  
none, 33  
nor, 9, 10, 43  
numbering, 26, 38  
“obtain that”, 15, 16, 47  
of, 25, 26, 29  
one, 23, 38  
only, 29, 42, 48  
ordinal numbers, 25, 34, 38  
  
paragraph, 4, 47  
participles, 30, 48  
  
percent, 35  
print, 8  
  
same, 36, 40  
satisfy, 48  
say, 11  
second largest, 36  
section, 4, 47  
shortly, 7, 48  
similar, 16, 48  
similarly, 48  
since, 15, 47  
smaller, 35  
smallest possible, 36  
so is, 10, 39  
some, 33  
succeed, 30  
such, 39, 47  
such that, 8  
  
that, 38  
the, 24  
the one, 38  
therefore, 15, 41, 45  
there is, 33  
these, 39  
thesis, 18, 20, 47  
the two, 34, 39  
this, 40  
this last, 39  
those, 38  
thus, 15, 41, 45  
to be defined, 9, 28  
too, 27  
to this end, 14, 47  
twice as long as, 36  
two-thirds, 35  
typefaces, 8  
  
unique, 24, 34  
union, 25  
unlikely, 28  
up to, 35, 38  
  
what, 15, 18, 47  
which, 15, 18, 47  
with, 26  
worth, 30  
worth while, 30